Patient Package Insert
Prescription Only Medicine
Carefully read the entire leaflet before starting use
ADCEF® Oral SUSPENSION & CAPSULES

COMPOSITION:
Adcef® 125 Oral Suspension: Each 5 ml contains 125 mg cefdinir.
Adcef® 250 Oral Suspension: Each 5 ml contains 250 mg cefdinir.
Adcef® 300 Capsules: Each capsule contains 300 mg cefdinir.
THERAPEUTIC ACTIVITY: Broad-spectrum antibiotic, semisynthetic cephalosporin.
APPROVED INDICATIONS: It is indicated in the treatment of infections caused by micro-organisms susceptible to cefdinir.
WHEN SHOULD THE PREPARATION NOT BE USED?

In patients with known hypersensitivity to the active ingredient, or to the cephalosporin group or to Penicillin's.

DO NOT TAKE THIS MEDICINE WITHOUT CONSULTING A PHYSICIAN:
- If you or your child is suffering or have suffered in the past from impaired kidney function, or if you or your child are under treatment of kidney failure using hemodialysis.
- If your child is diabetic, Adcef® oral suspension contains sugar; this could cause an increase in blood sugar levels.
WARNINGS: Do not take this medicine frequently or for long period for you or for your child without consulting a physician. If you or your child is allergic to any type of food or medicine, inform your physician before starting treatment with this medicine. You should tell your physician if you or your child suffers from diarrhea subsequent to the administration of antibacterial agents (colitis). This drug may give false result for glucose and ketone tests.
Note: This medicine contains sodium.
DRUG INTERACTIONS: If you or your child are taking or just finished taking any other prescriptions or over the counter medications, please notify your doctor or pharmacist, in order to prevent adverse reactions or lack of efficacy arising from drug interactions, this is especially important for medicines belonging to the following groups: Antacids that contain aluminium or magnesium, Iron supplements, Multivitamins containing iron, Probenecid (used to treat increased uric acid in blood).
• If you or your child take antacids, iron supplements, or multivitamins with iron, be sure to give Adcef® at least 2 hours before or 2 hours after the supplement.
• *Iron-fortified infant formula:* Adcef® may be taken with iron-fortified infant formula. Iron-fortified infant formula does not significantly interfere with drug absorption. The combination of iron and Adcef® sometimes turns the stool a brick red color, this reddish discoloration due to the combination of iron and Adcef® is usually not dangerous and not a cause for concern. If it does occur, red stool will stop when Adcef therapy is completed.
SIDE EFFECTS: In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine. Symptoms such as: diarrhea, rash and vomiting, vaginal moniliasis, abdominal pain, vaginitis, dyspepsia, flatulence, dry mouth. In the event that you experienced side effects not mentioned in this leaflet, or if there was a change in your general health, consult your doctor immediately.
- If you or your child experiences any of the following side effects, stop taking Adcef and contact your doctor immediately:
*** An allergic reaction (difficulty in breathing; swelling of the lips, face, or tongue; or a rash, redness, or itching).**
*** Episodes of nausea, vomiting, or diarrhea.**
*** Mucous or blood in the stool; or unusual bleeding.**
RECOMMENDED DOSAGE: The usual recommended dose is according to physician's instructions only. Do not exceed the recommended dosage. This medicine is to be taken at specific time intervals as determined by the physician.
MISSED DOSE: If a dose is missed, take the dose or give it to your child as soon as you remember but never take a double dose.
DIRECTIONS FOR USE: Adcef® suspension and capsules can be taken with or without food, but if stomach upset occurs take it with food.
Adcef® 125 & 250 mg Oral Suspension: Invert bottle and shake powder to become loose, and then add boiled and cooled water up to the line in two portions shaking after each addition to obtain 60 ml of homogeneous suspension. Shake well before each use. Do not take for more than 10 days after reconstitution. Take care to measure the dose using a plastic measuring spoon of 10 ml.
Adcef 300 Capsules: Do not chew! Or bite, or open the capsule. Capsules should be swallowed whole with a small amount of water.
HOW CAN YOU CONTRIBUTE TO THE SUCCESS OF THE TREATMENT? Complete the full course of treatment as instructed by the physician. Even if there is an improvement in your health, do not discontinue using this medicine, before consulting your physician.
AVOID POISONING: This medicine contains substances, which can be poisonous or fatal in large doses to children and/or infants, therefore must be kept out of their reach. This medicine has been prescribed for the treatment of your illness; it may harm other patients. Do not give this medicine to your relatives, neighbours, or anybody else. Do not take medicine in the dark! Check the label and the dose each time you take your medicine.
OVERDOSE: In case of accidental overdose or if a child takes this medicine by accident, seek professional assistance or rush to a hospital emergency room accompanied with the medicine's package. Do not induce vomiting unless explicitly instructed to do so by a doctor!
STORAGE: Store the capsules, the dry powder and reconstituted suspension at temperature not exceeding 25°C. Don't refrigerate. After reconstitution, the suspension may be used for 10 days, after which any unused portion must be discarded. The container should be kept tightly closed, and the suspension should be shaken well before each administration. Even if kept in their original container and stored as recommended; medicine may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt consult the pharmacist who dispensed the medicine to you.

PRESENTATION:
Adcef® 125 Oral Suspension: Brown glass bottle with powder for preparing 60 ml suspension with plastic measuring graduated spoon up to 10 ml.
Adcef® 250 Oral Suspension: Brown glass bottle with powder for preparing 60 ml suspension with plastic measuring graduated spoon up to 10 ml.
Adcef® 300 Capsules: A carton box with one milky plastic jar contains 10 & 14 capsules.

Pharmacare PLC- Beitunia- Industrial Zone- P.O. Box 677 - Ramallah, Palestine

L-2016122531/4418/3

نشرة للمريض
ملزم بوصفة طبية
يجب قراءة النشرة جيدا قبل البدء بالاستعمال
ادسف® معلق وكبسولات

التركيب الدوائي:
ادسف® ١٢٥ معلق: كل ٥ مل يحتوي على ١٢٥ ملغم من سيفدنير.
ادسف® ٢٥٠ معلق: كل ٥ مل يحتوي على ٢٥٠ ملغم من سيفدنير.
ادسف® ٣٠٠ كبسولات: كل كبسولة تحتوي على ٣٠٠ ملغم من سيفدنير.
الفعالية العلاجية: مضاد حيوي واسع الطيف، ينتمي إلى مجموعة السيفالوسبورين الشبه مصنعة.
دواعي الاستعمال: لمعالجة الالتهابات الناجمة عن الجراثيم الحساسة للسيفدنير.
متى لا يجوز استعمال هذا الدواء؟

للمرضى المعروف ان لديهم حساسية للمادة الفعالة او لمجموعة السيفالوسبورين او البنسيلين.

لا يجوز استعمال هذا الدواء قبل استشارة الطبيب: إذا كنت تعاني أنت أو طفلك أو عانيت في الماضي من خلل في وظيفة الكلى، أو إذا كنت أنت أو طفلك خاضعين لعلاج الفشل الكلوي بديلزة الدم. إذا كان طفلك مصاب بداء السكري، فان الادسف® معلق يحتوي على السكر، وهذا يؤدي إلى زيادة نسبة السكر في الدم.
التحذيرات: يجب تجنب العلاج الطويل الأمد والمستمر لك ولطفلك دون استشارة الطبيب. يجب إبلاغ الطبيب قبل استعمال هذا الدواء إن تواجدت حالات الحساسية لدواء معين او طعام ما لديك او لدى طفلك. يجب إخبار الطبيب إذا كنت تعاني أو إذا كان طفلك يعاني من إسهال نتيجة لاستخدام المضادات الحيوية (التهاب القولون). تناول هذا الدواء قد يؤدي إلى إمكانية إعطاء نتيجة خاطئة لفحصي الجلوكوز والكيتون.
ملاحظة: يحتوي هذا المستحضر على الصوديوم.
التداخلات الدوائية: إذا كنت أنت أو طفلك تتناولون أو أنهيتم قبل فترة وجيزة تناول أية أدوية أخرى بما فيها الأدوية التي تصرف بدون وصفة طبية، اعلم طبيبك المعالج أو الصيدلاني وذلك لمنع التفاعلات الجانبية أو عدم النجاعة الناجمة عن التفاعلات ما بين الأدوية وهذا مهم خاصة مع الأدوية التي تنتمي الى المجموعات التالية: مضادات الحموضة التي تحتوي على الألمنيوم أو المغنيسيوم، الحديد ومشتقاته، الفيتامينات التي تحتوي على الحديد، البروبنسيد (لعلاج فرط حامض البول في الدم).
• إذا كنت أو إذا كان طفلك يتناول مضادات الحموضة، أو الحديد ومشتقاته، أو الفيتامينات التي تحتوي على الحديد: يجب إعطاء الادسف® بعد ساعتين أو قبل ساعتين على الاقل من تناوله لأحد هذه الأدوية.
• الحديد المدعم للأطفال: يمكن إعطاء الادسف® لطفلك مع تركيبة الحديد المدعم مثل حليب الأطفال المدعم بالحديد، لان الحديد المدعم الموجود في هذه التركيبات لا يتداخل بشكل ملحوظ مع امتصاص الادسف® في الجسم، ولكن تداخل الادسف® مع الحديد يعطي في بعض الأحيان اللون الأحمر القرميدي للبراز والذي لا يعد خطير وان حصل فانه يزول عادة بعد إكمال العلاج بالادسف®.
الأعراض الجانبية: بالإضافة إلى التأثير الطبي للدواء فان لكل دواء تأثيرات وأعراض جانبية غير مرغوب فيها قد تحدث أثناء تناول هذا الدواء. مثال ذلك: الإسهال، الطفح الجلدي، القئ، داء الطوقيات المهبلي، آلام في البطن، التهاب مهبلي، تخمة، انتفاخ في البطن، جفاف في الفم. أما إذا ظهرت لديك أعراض جانبية لم تذكر في هذه النشرة، أو إذا طرأ أي تغيير على شعورك العام نتيجة لتناولك لهذا الدواء، عليك عندئذ مراجعة الطبيب حالا.
إذا عانيت أو عانى طفلك من أي من الأعراض الجانبية التالية يجب أن تتوقف عن تناول هذا الدواء وان تراجع طبيبك في الحال:
* تفاعلات الحساسية (صعوبة في التنفس، انتفاخ في الشفتين، الوجه، أو اللسان أو ظهور الطفح الجلدي، الاحمرار أو الحكة).
* نوبات غثيان، قئ، أو إسهال.
* ظهور المخاط أو الدم في البراز، أو أي نزيف غير عادي.
الجرعة الدوائية: الجرعة الموصى بها تكون حسب إرشادات الطبيب فقط. يجب عدم تجاوز الجرعة المحددة. يجب تناول هذا الدواء في فترات معينة كما حددها الطبيب.
الجرعة المنسية: في حالة نسيان تناول الدواء في الوقت المناسب، تناول الدواء او أعطه لطفلك فور تذكرك. ولكن لا يجب تناول جرعة مضاعفة.
إرشادات الاستعمال: يمكن تناول ادسف® كبسولات ومعلق مع أو بدون طعام، ولكن إذا حدثت اضطرا بات في المعدة جراء ذلك، يمكن تناوله مع الطعام.
ادسف® ١٢٥و٢٥٠ ملغم معلق: اقلب الزجاجة لكي تحرك البودرة داخلها ثم أضف كمية كافية من الماء المغلي والمبرد على دفعتين مع الخض حتى يصل حجم المعلق إلى الخط المرسوم على العبوة للحصول على ٦٠ مل من المعلق المتجانس. خض الزجاجة جيدا قبل كل استعمال. لا تتناول الدواء بعد ١٠ أيام من تحضيره. احرص على قياس الجرعة باستخدام ملعقة القياس البلاستيكية المرفقة بالعبوة والمدرجة الى ١٠ مل.
ادسف® ٣٠٠ كبسولات: لا تمضغ! أو تقضم أو تفتح الكبسولة. ابلع الكبسولة كاملة مع قليل من الماء.
كيف يمكنك المساهمة في إنجاح العلاج: إكمال دورة العلاج بشكل كامل كما هو مقرر من قبل الطبيب. عدم التوقف عن تناول العلاج دون استشارة الطبيب حتى لو طرأ تحسن على الصحة العامة.
لتجنب التسمم: هذا الدواء يحتوي على مواد قد تكون سامة أو قاتلة للأطفال أو الرضع بجرعات عالية، لذلك يجب حفظ الأدوية في مكان بعيد عن متناول أيدي الأطفال أو الرضع وذلك لتفادي إصابتهم بالتسمم. هذا الدواء وصف لعلاج مرضك أنت لذا قد يسبب الضرر لمريض آخر. لا تعط هذا الدواء لجارك أو قريبك أو أي شخص آخر. لا يجوز تناول هذا الدواء في الظلام. يجب التأكد من الدواء و الجرعة قبل تناولك للدواء.
الجرعة الزائدة: في حال تناول جرعة زائدة أو إذا قام الطفل بتناول هذا الدواء بالخطأ يجب التوجه إلى الطوارئ مصطحبا معك علبة الدواء. لا تسبب التقيؤ بدون تعليمات صريحة من الطبيب.

التخزين: تحفظ الكبسولات والبودرة الجافة والمعلق بعد تحضيره في درجة حرارة لا تتجاوز ٢٥ درجة مئوية. لا يحفظ هذا الدواء في الثلاجة. بعد التحضير يبقى المعلق صالحا للاستعمال لمدة ١٠ ايام، ويجب اتلاف اي كمية غير مستعملة من المعلق بعد هذه المدة. يجب الحفاظ على اغلاق العبوة جيدا بعد تناولك لهذا الدواء، كما ويجب خض العبوة جيدا قبل كل استعمال. لا تبقى الأدوية صالحة لفترات طويلة حتى ولو ظلت في علبتها الأصلية وتم حفظها بموجب الإرشادات. الرجاء ملاحظة تاريخ انتهاء صلاحية المستحضر. في حالة الشك يجب استشارة الصيدلي الذي صرف لك الدواء.

الشكل الصيدلاني والمحتوى:
ادسف® ١٢٥ معلق: قنينة زجاجية بنية اللون مع بودرة لتحضير معلق (٦٠ مل) مع ملعقة بلاستيكية مدرجة الى ١٠ مل.
ادسف® ٢٥٠ معلق: قنينة زجاجية بنية اللون مع بودرة لتحضير معلق (٦٠ مل) مع ملعقة بلاستيكية مدرجة الى ١٠ مل.
ادسف® ٣٠٠ كبسولات: كرتونة تحتوي على عبوة بلاستيكية بيضاء تحتوي على ١٠ و ١٤ كبسولة.

شركة دار الشفاء لصناعة الأدوية- بيتونيا – المنطقة الصناعية- ص.ب. ٦٧٧- رام الله – فلسطين
٣/٤٤١٨/٢٠١٦١٢٢٥٣١-ل